



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

B 447300



Library of the University of Michigan

*Bought with the income
of the*

*Ford - Messer
Bequest*



E. F. PARKER

Science Library

QK

607. -

.C782

ILLUSTRATIONS
OF
BRITISH FUNGI.

ILLUSTRATIONS
OF
BRITISH FUNGI
(HYMENOMYCETES),

To serve as an Atlas to the "Handbook of British Fungi."

BY

*Morke's
dubitt*
M. C. COOKE, M.A., LL.D., A.L.S.,

President of the Quekett Microscopical Club, and of the Hackney Natural History and Microscopical Society. Fellow of the Botanical Society of Edinburgh. Corresponding Member of the Academy of Natural Sciences, Philadelphia; of the Academy of Science of New York; of the Portland Society of Natural History, Maine; and of the Società Crittogamologica Italiana. Honorary Member of the Cryptogamic Society of Scotland; the Woolhope Field Club; the Hertfordshire Natural History Society; the Norfolk and Norwich Naturalists' Society; the Essex Field Club; the Birmingham Natural History Society; and other Societies

VOL. VII.

London:

WILLIAMS AND NORGATE, HENRIETTA STREET, W.C.

1888-1890.

SYSTEMATIC INDEX.

GEN. 9. **LACTARIUS**, *Fr. Epic.* 333.

Hymenophore continuous with the stem. Gills unequal, between membranaceous and waxy, rather rigid, *milky*, edge acute. Spores globose, white, rarely turning yellowish. Fleshy, terrestrial, putrescent fungi, with a depressed pileus, and adnate-decurrent, sometimes branched gills.

Tribe 1. Piperites. Stem central. Gills unchangeable, naked, not discoloured or pruinose. Milk at first white, commonly acid.

* **TRICHOLOMOIDEI.** Pileus moist, viscid, margin at first involute and tomentose.

922.	scrobiculatus, <i>Scop.</i>	.	.	971
923.	torminosus, <i>Schæff.</i>	.	.	972
924.	cilicioides, <i>Fries</i>	.	.	973
—	intermedius, <i>Kromb.</i>	.	.	—
925.	turpis, <i>Weinm.</i>	.	.	987
—	var. plumbeus, <i>Fries.</i>	.	.	—
926.	controversus, <i>Pers.</i>	.	.	1003
927.	pubescens, <i>Fries</i>	.	.	974
928.	aspideus, <i>Fries</i>	.	.	1083

** **LIMACINI.** Pileus moist, viscid, with a pellicle; margin naked.

929.	insulsus, <i>Fries</i>	.	.	975
—	zonarius, <i>Bull</i>	.	.	—
930.	utilis, <i>Weinm.</i>	.	.	1084
931.	blennius, <i>Fries</i>	.	.	988
932.	hyginus, <i>Fries</i>	.	.	989
933.	trivialis, <i>Fries</i>	.	.	976
934.	circellatus, <i>Batt.</i>	.	.	990
935.	avidus, <i>Fries</i>	.	.	991

*** **PIPERATI.** Pileus without pellicle, hence absolutely dry, often unpolished.

936.	flexuosus, <i>Fries</i>	.	.	992
937.	pyrogalus, <i>Bull</i>	.	.	993
938.	squalidus, <i>Kromb.</i>	.	.	1004a
939.	capsicum, <i>Schulz.</i>	.	.	977
940.	chrysorrheus, <i>Fries</i>	.	.	984
941.	acris, <i>Bolt.</i>	.	.	1005

Dec 4 - 13-37 2014

942.	umbrinus, <i>Pers.</i>	.	.	1006
943.	pargamenus, <i>Swartz.</i>	.	.	978
944.	piperatus, <i>Scop.</i>	.	.	979
945.	vellerens, <i>Fries</i>	.	.	980
946.	exsuccus, <i>Otto.</i>	.	.	981
938.	scoticus, <i>B. & Br.</i> ? (<i>1892</i>)	.	.	1004b

Tribe 2. Dapetes. Stem central. Gills naked, milk at first brightly coloured.

947.	deliciosus, <i>Linn.</i>	.	.	982
------	--------------------------	---	---	-----

Tribe 3. Russularia. Stem central. Gills pallid, then discoloured, then darker, changing with the light, at length pruinose with white. Milk at first white, either mild, or mild then becoming acrid.

* *Pileus at first viscid.*

948.	pallidus, <i>Pers.</i>	.	.	1007
949.	quietus, <i>Fries</i>	.	.	983
950.	aurantiacus, <i>Fl. Dan.</i>	.	.	1099
—	theiogalus, <i>Bull</i>	.	.	—
951.	cremor, <i>Fries</i>	}	.	1008
	var. pauper, <i>Karst.</i>			
952.	vietus, <i>Fries</i>	.	.	1009
953.	cyathula, <i>Fries</i>	.	.	1085

** *Pileus unpolished, squamulose, villous, or pruinose.*

954.	rufus, <i>Scop.</i>	.	.	985
955.	helvus, <i>Fries</i>	.	.	994
956.	tomentosus, <i>Otto.</i>	.	.	1010
—	mammosus, <i>Fries</i>	.	.	—
957.	var. monstrosus, <i>Fr.</i>	.	.	995
958.	glyciosmus, <i>Fries</i>	.	.	1011
959.	fuliginosus, <i>Fries</i>	.	.	996
960.	picinus, <i>Fries</i>	.	.	997
961.	{ lilacinus, <i>Lasch.</i>	.	.	998a
	{ spinosulus, <i>Quel.</i> var.	.	.	998b

*** *Pileus polished, smooth.*

962.	volemus, <i>Fries</i>	.	.	999
963.	ichoratus, <i>Batsch.</i>	.	.	1000
964.	serifluus, <i>D. Cand.</i>	.	.	1012
965.	mitissimus, <i>Fries</i>	.	.	1001
966.	subdulcis, <i>Bull</i>	.	.	1002
—	var. Terrei, <i>B & Br.</i>	.	.	—
967.	{ camphoratus, <i>Fries</i>	.	.	1013a
	{ cimicarius, <i>Batsch.</i>	.	.	1013b
968a.	subumbonatus, <i>Lind.</i>	.	.	986a
969a.	obnubilis, <i>Lasch.</i>	.	.	1014a
968b.	minimus, <i>Smith</i>	.	.	986b

Tribe 4. *Pleuropus*. Stem excentric or lateral,

962 969b. obliquus, *Fries*. *Muth* : 7a . 1014b = 1-filic

GEN. 10. **RUSSULA**, *Pers*

Veil none. Hymenophore descending unchanged into the vesiculose trama. Gills rigid, fragile, not milky, edge acute. Spores rounded, often echinulate, white, or becoming yellow. Terrestrial, fleshy, putrescent fungi, stem polished, pileus at first or at length depressed.

- I. **COMPACTÆ**. Pileus everywhere fleshy, hence the margin at first turned in, always without striæ, or a distinct viscid pellicle (hence the colour not variable, only changing with age or the action of the air), flesh firm, compact. Stem solid, fleshy. Gills unequal.

970.	<i>nigricans</i> , <i>Bull.</i>	.	1015
971.	var. <i>albo-nigra</i> , <i>Krom.</i>	.	1016
972.	<i>adusta</i> , <i>Pers.</i>	.	1051
973.	<i>densifolia</i> , <i>Secr.</i>	.	1017
974.	<i>semicrema</i> , <i>Fries</i>	.	1067
975.	<i>delica</i> , <i>Fries</i>	.	1068
—	<i>elephantina</i> , <i>Fries</i>	.	?
976.	<i>mustelina</i> , <i>Fries</i>	.	1018

- II. **FURCATÆ**. Pileus compact, firm, covered with a thin, closely adnate pellicle, which at length disappears, margin abruptly thin, at first turned in, then spreading, acute, even. Stem at first compact, at length soft and spongy within. Gills rather forked, mixed with a few shorter ones, commonly attenuated towards each end, thin and normally narrow.

977.	<i>olivascens</i> , <i>Fries</i>	.	1035
978.	<i>furcata</i> , <i>Pers.</i>	.	1036
979.	var. <i>pictipes</i> , <i>Cke.</i>	.	1086
980.	var. <i>ochroviridis</i> , <i>Cke.</i>	.	1100
981.	<i>sanguinea</i> , <i>Bull</i>	.	1019
982.	<i>rosacea</i> , <i>Fries</i>	.	1020
983.	<i>maculata</i> , <i>Quel.</i>	.	1069
984.	<i>sardonia</i> , <i>Fries</i>	.	1037
985.	<i>depallens</i> , <i>Fries</i>	.	1021
986.	<i>purpurea</i> , <i>Gillet</i>	.	1022
987.	<i>cærulea</i> , <i>Pers.</i>	.	1052
988.	<i>drimæia</i> , <i>Cooke</i>	.	1023

- III. **RIGIDÆ**. Pileus destitute of a viscid pellicle, absolutely dry, rigid, the cuticle commonly breaking up into flocci or granules. Flesh

thick, compact, firm, vanishing near the margin, which is straight (never involute), spreading, and without striæ. Stem solid, at first hard, then softer and spongy. Of the gills, a few reach half-way to the stem; the others are divided, rigid, dilated in front, and extend into a very broad, rounded end, whence the margin of the pileus becomes obtuse and not inflexed.

989.	<i>lactea, Pers.</i>	.	.	1070
990.	var. <i>incarnata</i>	.	.	1071
991.	<i>virescens, Schæff.</i>	.	.	1039
992, 993.	<i>cutefracta, Cooke</i>	.	1024, 1040	
994, 995.	<i>lepida, Fries</i>	.	1072, 1073	
996.	<i>rubra, D. Cand.</i>	.	.	1025
997.	var. <i>atropurpurea, Kromb.</i>	.	.	1087
998.	<i>Linnæi, Fries</i>	.	.	1026
999, 1000.	<i>xerampelina, Schæff.</i>	.	1053, 1074	
1001.	<i>olivacea, Schæff.</i>	.	.	1041

IV. HETEROPHYLLÆ. Pileus fleshy, firm, covered with a thin adnate pellicle, margin thin, at first turned in, then expanded and striate. Gills many shorter mixed with the longer ones, others forked. Stem solid, stout, spongy within.

1002.	<i>vesca, Fries</i>	.	.	1075
1003b.	<i>Du Portii, Phil.</i>	.	.	1042b
1004.	<i>lilacea, Quelet</i>	.	.	1054
1005.	<i>azurea, Bres.</i>	.	.	1088
1003a.	<i>serotina, Quelet</i>	.	.	1042a
1006, 1007, 1008.	<i>cyanozantha, Schæff.</i>	1043, 1076, 1077		
1009, 1010.	<i>heterophylla, Fries</i>	1044, 1045		
1011.	<i>galochroa, Bull</i>	.	.	1089
1012.	<i>consobrina, Fries</i>	.	.	1055
1013.	var. <i>intermedia, Cke.</i>	.	.	1056
1014.	var. <i>sororia, Larb.</i>	.	.	1057
1015.	<i>foetens, Pers.</i>	.	.	1046
1016.	<i>subfoetens, Smith</i>	.	.	1047
1017.	<i>fellea, Fries</i>	.	.	1058
1018.	<i>elegans, Bres.</i>	.	.	1027
1019.	<i>Quelettii, Fries</i>	.	.	1028
1020.	<i>expallens, Gillet</i>	.	.	1029

V. FRAGILES. Pileus more or less fleshy, rigid, but fragile, covered with a pellicle, always continuous, viscid after rain, and somewhat separable, margin membranaceous, at first closed in, but not involute, when adult commonly sulcate and

tuberculose. Flesh usually floccose, lax and friable. Stem spongy, at length soft and hollow. Gills almost all equal, simple, broadest in front, free in the pileus when closed.

* Gills and spores white.

1021.	<i>cmetica</i> , <i>Fries</i>	.	.	1030
1022.	var. <i>clusii</i> , <i>Vitt.</i>	.	.	1031
1023.	var. <i>fallax</i> , <i>Schæff.</i>	.	.	1059
1024.	<i>pectinata</i> , <i>Bull.</i>	.	.	1101
1025.	<i>ochroleuca</i> , <i>Pers.</i>	.	.	1049
1026.	<i>granulosa</i> , <i>Cooke</i>	.	.	1038
1027.	<i>æruginea</i> , <i>Fries</i>	.	.	1090
1028.	<i>fragilis</i> , <i>Pers.</i>	.	.	1091
1029.	{ var. <i>violacea</i> , <i>Fr.</i>	.	.	1060a
	{ var. <i>nivea</i> , <i>Fr.</i>	.	.	1060b
1030.	<i>figibilis</i> , <i>Britz.</i>	.	.	1048
1031.	<i>citrina</i> , <i>Gillet</i>	.	.	1078
1032.	<i>punctata</i> , <i>Gillet</i>	.	.	1082

** Gills and spores white, then yellow or bright lemon.

1033, 1034.	<i>veternosa</i> , <i>Fries</i>	.	1033, 1092
1035.	var. <i>roseipes</i> , <i>Bres.</i>	.	1081
1036, 1037.	<i>integra</i> , <i>Linn.</i>	.	1084, 1093
1038.	var. <i>alba</i> , <i>Cke.</i>	.	1094
1039.	<i>decolorans</i> , <i>Fries</i>	.	1079
1040.	<i>Barlx</i> , <i>Quelet.</i>	.	1061
1041.	<i>aurata</i> , <i>With.</i>	.	1080
1042, 1043.	<i>nitida</i> , <i>Pers.</i>	.	1062, 1063
1044.	{ var. <i>cuprea</i> , <i>Krom.</i>	.	1095b
	{ <i>pulchralis</i> , <i>Britz.</i>	.	1095a
1045.	<i>armeniaca</i> , <i>Cooke</i>	.	1064
1046, 1047.	<i>puellaris</i> , <i>Fries</i>	.	1065, 1066

. Gills and spores ochre.

1048, 1049.	<i>alutacea</i> , <i>Fries</i>	.	1096, 1097
1050.	<i>ochracea</i> , <i>A. & S.</i>	.	1050
1051.	<i>lutea</i> , <i>Huds.</i>	.	1082
1052.	<i>nauseosa</i> , <i>Pers.</i> (<i>Alut. 1102a</i>)	.	1147
1053.	<i>vitellina</i> , <i>Pers.</i>	.	1102b
1054.	<i>chameleontina</i> , <i>Fries</i>	.	1098

GEN. 11. **CANTHARELLUS**, *Adans.*

Hymenophore continuous with the stem, descending unchanged into the trama. Gills thick, between fleshy and waxy, fold-like, rather branched, edge obtuse. Spores white. Fleshy membranaceous putrescent fungi, veil none.

1. **MESOPUS**. Pileus entire. Stem central.

* *Pileus and solid stem fleshy.*

1055.	<i>cibarius, Fries</i>	.	1103
1056.	{ <i>var. rufipes</i>	.	1131a
	<i>Friesii, Quelet.</i>	.	1131b
1057.	<i>aurantiacus, Wulf.</i>	.	1104
1058a.	<i>Brownii, B. & Br.</i>	.	1106a
1059.	<i>carbonarius, A. & S.</i>	.	1105
	<i>var. radicosus, B. & Br.</i>	}	1105
1058b.	<i>umbonatus, Fries</i>	.	1106b
1060a.	<i>albidus, Fries</i>	.	1107a

** *Pileus submembranaceous, stem hollow, polished.*

1060b.	<i>Houghtoni, Phil.</i>	.	1107b
1061.	<i>tubæformis, Fries</i>	.	1108
	<i>var. lutescens, Bull.</i>	.	1109
1062.	<i>infundibuliformis, Fries</i>	.	1110a
1063a.	<i>cinereus, Pers.</i>	.	1110a
1064a.	<i>leucophæus, Nouel.</i>	.	1111a
1063b.	<i>cupulatus, Fries</i>	.	1110b
1098a.	<i>devevus, Fries</i>	.	1150a
1064b.	<i>Stevensoni, B. & Br.</i>	.	1111b

II. *PLEUROPUS.* *Dimidiate, stem lateral.*

1065.	{ <i>muscigenus, Bull.</i>	.	1115a
	<i>glaucus, Fries</i>	.	1115b

III. *RESUPINATI.* *Pileus entire, at first cup-shaped, fixed at the vertex, then subreflexed.** *Bryophili.*

1066.	{ <i>retirugus, Bull.</i>	.	1112a
	<i>lobatus, Fries</i>	.	1112b

GEN. 12. *NYCTALIS*, *Fr. Syst. Orb.* 208.

Hymenophore continuous with the stem. Gills fleshy, thick, juicy, edge obtuse, not decurrent on the stem nor fold-like. Veil floccosely pruinose. Fleshy fungi, not reviving, of uncertain occurrence.

I. *SPELEÆ.* *Gills crowded, somewhat growing together.*

1067a.	<i>caliginosa, Smith</i>	.	1132a
--------	--------------------------	---	-------

II. *PARASITÆ.* *Gills distinct, distant, growing on putrid fungi.*

1067b.	<i>asterophora, Fries</i>	.	1132b
1068.	<i>parasitica, Bull.</i>	.	1113

GEN. 13. *MARASMIUS*, *Fr. Gen. Hym.* 372.

Tough, dry, shrivelling (not putrescent), fungi reviving when moistened. Hymenophore continuous with the stem,

but heterogeneous, descending into the trama; veil none. Stem cartilaginous or horny. Gills tough, rather distant, edge acute and entire.

- I. COLLYBIA. Pileus fleshy, tough, at length rather leathery, sulcate or wrinkled, margin at first turned in. Stem somewhat cartilaginous. Mycelium floccose, or in a few not manifest.
- A. SCORTEI. Stem solid, or stuffed, with a pith, then hollow, fibrous within, externally with a villous down, easily wiped away, which covers the cartilaginous cuticle. Gills receding, free.
- * Stem woolly or strigose at the base.
- | | | | | |
|-------|-------------------------|---|---|------|
| 1069. | urens, <i>Bull.</i> | . | . | 1116 |
| 1070. | peronatus, <i>Bolt.</i> | . | . | 1117 |
| 1071. | porreus, <i>Pers.</i> | . | . | 1188 |
- ** Stem naked at the base, often interwoven with twisted fibres.
- | | | | | |
|-------|--------------------------|---|---|-------|
| 1072. | oreades, <i>Bolt.</i> | . | . | 1118 |
| 1073. | { plancus, <i>Fries</i> | . | . | 1119a |
| | { scorteus, <i>Fries</i> | . | . | 1119b |
- B. TERGINI. Stem rooting, definitely tubular, not fibrous, manifestly cartilaginous. Gills receding, free. Pileus thinner than in the preceding, hygrophanous, smooth, or striate at the margin.
- * Stem woolly below, smooth above.
- | | | | | |
|--------|--------------------------------|---|---|-------|
| 1074. | prasiosmus, <i>Fries</i> | . | . | 1120 |
| 1075. | { varicosus, <i>Fries</i> | . | . | 1121a |
| | { fuscopurpureus, <i>Pers.</i> | . | . | 1121b |
| 1076a. | terginus, <i>Fries</i> | . | . | 1122a |
- ** Stem everywhere pruinose, velvety.
- | | | | | |
|--------|-------------------------------|---|---|-------|
| 1077. | { Wynnei, <i>B. & Br.</i> | . | . | 1123a |
| | { erythropus, <i>Fries</i> | . | . | 1123b |
| 1076b. | archyropus, <i>Pers.</i> | . | . | 1122b |
| 1078. | { torquescens, <i>Quel.</i> | . | . | 1124a |
| | { impudicus, <i>Fries</i> | . | . | 1124b |
- C. CALOPODES. Stem short, not rooting, grafted, often with a floccose tubercle at the base. Pileus convex, involute, then plane and depressed, in which state the gills, typically adnate, are somewhat decurrent.
- * Stem very smooth and shining above, simple at the base.
- | | | | | |
|-------|---------------------------|---|---|-------|
| 1079. | { scorodoni, <i>Fries</i> | . | . | 1125a |
| | { calopus, <i>Pers.</i> | . | . | 1125b |

- | | | | | | |
|-------|---|----------------------------------|---|---|-------|
| | { | <i>vaillantii</i> , <i>Fries</i> | . | . | 1126a |
| 1080. | | <i>angulatus</i> , <i>Pers.</i> | . | . | 1126b |
| | } | <i>languidus</i> , <i>Fries</i> | . | . | 1126c |

** Stem velvety or pruinose, base rather tuberculose.

- | | | | | | |
|--------|---|---------------------------------|---|---|-------|
| 1081a. | | <i>foetidus</i> , <i>Sow.</i> | . | . | 1134a |
| 1082a. | | <i>amadelpus</i> , <i>Fries</i> | . | . | 1127a |
| 1082. | { | <i>ramealis</i> , <i>Bull.</i> | . | . | 1127b |
| | } | <i>candidus</i> , <i>Bolt.</i> | . | . | 1127c |

Tribe 2. Mycena. Stem horny, fistulose, sometimes medullate, tough, dry, mycelium rhizomorphoid, corticate not floccose. Pileus rather membranaceous, campanulate, then expanded, margin at first straight, pressed to the stem.

A. CHORDALES. Stem rigid, rooting, or adnate by a dilated base. Pileus campanulate or convex.

- | | | | | | |
|--------|---|------------------------------------|---|---|-------|
| 1083. | { | <i>alliaceus</i> , <i>Jacq.</i> | . | . | 1128a |
| | | <i>cohaerens</i> , <i>Pers.</i> | . | . | 1128b |
| 1081b. | | <i>cauticinalis</i> , <i>With.</i> | . | . | 1134b |

B. ROTULÆ. Stem filiform, flaccid, grafted at the base. Pileus soon rather plane or umbilicate.

* Stem quite smooth, shining.

- | | | | | | |
|-------|---|--------------------------------------|---|---|-------|
| 1084. | { | <i>rotula</i> , <i>Scop.</i> | . | . | 1129a |
| | | <i>graminum</i> , <i>Lib.</i> | . | . | 1129b |
| | | <i>androsaceus</i> , <i>Linn</i> | . | . | 1129c |
| 1085. | { | <i>splachnoides</i> , <i>Fries</i> | . | . | 1130a |
| | } | <i>Curreyi</i> , <i>B. & Br.</i> | . | . | 1130b |

** Stem velvety or pilose.

- | | | | | | |
|--------|---|---|---|---|-------|
| 1085c. | | <i>perforans</i> , <i>Fries</i> | . | . | 1130c |
| 1086. | { | <i>institius</i> , <i>Fries</i> | . | . | 1135a |
| | | <i>Hudsoni</i> , <i>Pers.</i> | . | . | 1135b |
| | | <i>epichlœe</i> , <i>Fries</i> | . | . | 1136a |
| 1087. | { | <i>actinophorus</i> , <i>B. & Br.</i> | . | . | 1136b |
| | | <i>saccharinus</i> , <i>Fries</i> | . | . | 1136c |
| 1088. | { | <i>epiphyllus</i> , <i>Fries</i> | . | . | 1137a |
| | } | <i>polyadelphus</i> , <i>Lasch.</i> | . | . | 1137b |

Tribe 3. Apus. Pileus sessile, resupinate.

- | | | | | |
|--------|--|---|---|---------|
| 1088c. | <i>spodoleucus</i> , <i>Berk.</i> | . | . | } 1137c |
| | including <i>M Broomei</i> , <i>B.</i> | . | . | |

GEN. 14. LENTINUS, *Fries.*

Pileus between fleshy and leathery, tough, hard when old, persistent. Hymenophore continuous with the stem, or base of the pileus when sessile. Gills concrete with the

hymenophore, thin, unequal, membranaceous, with the edge serrate or torn into teeth. Spores white, roundish, even. Growing on wood.

I. **MESOPODES.** Pileus nearly entire, stem distinct.

* *Lepidei.* Pileus squamose, more or less manifestly veiled.

1089.	tigrinus, <i>Bull.</i>	.	1138, 1139	
1090.	Dunalii, <i>D. Cand.</i>	.	1139	
1091, 1092.	lepideus, <i>Fries</i>	.	1140, 1141	

** *Pulverulenti.* Pileus villose or pulverulent.

---	leontopodius, <i>Schultz</i>	.	---	
---	pulverulentus, <i>Scop.</i>	.	---	
---	resinaceus, <i>Trog.</i>	.	---	
---	adhærens, <i>Alb. & Schw.</i>	.	---	

*** *Cochleati.* Pileus smooth.

1093a.	cochleatus, <i>Fries</i>	.	1142a	
--------	--------------------------	---	-------	--

II. **PLEUROT.** Dimidiate, sessile, or with a rather lateral stem.

1094.	scoticus, <i>B. & Br.</i>	.	1143	
1095a.	fimbriatus, <i>Curr.</i>	.	1148a	
1093b.	vulpinus, <i>Fries</i>	.	1142b	
1095b.	flabelliformis, <i>Bolt.</i>	.	1148b	

GEN. 15. **PANUS**, *Fries Epicr.* 396.

Whole fungus fleshy-leathery, tough, becoming dry, substance fibrous, radiating into the hymenium. Gills concrete with the hymenophore, unequal, at length leathery, edge entire. Spores even, white.

* Pileus irregular, stem excentric.

1096.	{ conchatus, <i>Fries</i>	.	1149a	
	{ torulosus, <i>Fries</i>	.	1149b	

** Stem definitely lateral.

1097.	{ stypticus, <i>Bull.</i>	.	1144a	
	{ farinaceus, <i>Schum.</i>	.	1144b	

*** Pileus resupinate, sessile, or extended behind.

1097c.	patellaris, <i>Fries</i>	.	1144c	
---	Stevensoni, <i>B. & Br.</i>	.	---	

GEN. 16. **XEROTUS**, *Fries Elen.* I. 48.

Hymenophore continuous with the stem, descending into the trama, which is similar to the leathery pileus. Gills leathery, broadly fold-like, dichotomous, edge entire, obtuse.

1098 =

1098 - (mispl. page 6)

1089b. degener, *Fries* . . . 1150bGEN. 17. **TROGIA**, *Fries Epic.* 402.

Gills fold-like, longitudinally channelled along the edge,
in other respects like *Xerotus*.

1099a crispa, *Pers.* . . . 1114aGEN. 18. **SCHIZOPHYLLUM**, *Fries Obs.* 1. 103.

Pileus without flesh, dry. Gills leathery, branched in a
fan-like manner, united above by the tomentose pellicle, bifid,
edge longitudinally split. Spores roundish, white.

1099b commune, *Fries* . . . 1114bGEN. 19. **LENZITES**, *Fries Epic.* 403.

Pileus corky or leathery, substance dry and floccose. Gills
leathery, firm, either simple, unequal, or anastomosing and
forming pores behind, trama floccose and similar to the
pileus, edge rather acute.

* Growing on deciduous trees.

1100. { *betulina*, *Linn* . . . 1145a
 { *flaccida*, *Bull.* . . . 1145b

** Growing on conifers.

1101. { *sepiaria*, *Fries* . . . 1146a
 { *abietina*, *Bull.* . . . 1146b

100%

87

No 922

PL. 971.



LACTARIUS (PIPERITES) SCROBICULATUS. Scop.
on the ground, near Bristol.

۱۲۰۵

York

W40U



LACTARIUS (PIPERITES) TORMINOSUS. *Schaff.*
on the ground in woods. Aymestry, Oct. 1871.

1000

4401

1901

No 924

PL 973.



LACTARIUS (PIPERITES) CILICIOIDES. *Fries.*
on the ground. near Hereford.



1901

Ver 34

16 4

No 925

PL. 987.

milk white, acrid.



viscid.

8 μ

LACTARIUS (PIPERITES) TURPIS. Weinm.
in woods. Dinmore, Oct. 1881.

৪৭০১

1074

1701

milk white, acrid.



6-8/M

LACTARIUS (PIPERITES) CONTROVERSUS. Pers.
in pastures, &c. near Hereford.

1490

UorM

No 927

PL 974.



LACTARIUS (PIPERITES) PUBESCENS. *Fries.*
in open grassy places. Ledbury, Oct. 1888.

1701

1701

1701


1701

1074

870

No 928

PL. 1083.



LACTARIUS (PIPERITES) ASPIDEUS. *Fries.*
in swampy places. Harewood, near Leeds. Sept. 1888.

Mr. U

Udemy

1870

40429

PL 978.



LACTARIUS (PIPERATES) INSULSUS. *Fries.*
in woods. Mattishall, Oct. 1884.

8701



1871

WFOU



No 930

PL. 1084.



LACTARIUS (PIPERITES) UTILIS. Weism.
on the ground. near Grove Park, Warwick, 1888.

Mr. U



U of M

No 931

PL. 988.



LACTARIUS (PIPERITES) BLENNIUS. Fries.
in beech woods. Epping, 1884.

Mr. U

U of M

88

No 932

PL. 989.



LACTARIUS (PIPERITES) HYGINUS. *Fries.*
amongst grass. Letton Court and Epping Forest, 1886.

M70U

UofM

177

20 923

PL. 976.

milk white, acrid.



LACTARIUS (PIPERITES) TRIVIALIS. *Fries.*
in moist woods. near Hereford.

Wm. W. W.

Uor 34

84

No 934

PL. 990.



LACTARIUS (PIPERITES) CIRCELLATUS. *Fries.*
amongst grass. Highgate. Oct. 1883.

1700

UofA

1000

260955

PL. 991.



LACTARIUS (PIPERITES) UVIDUS. *Fries.*
in moist woods. Loughton. Aug. 1872.

www

Mr. U

No 936

PL. 992.

milk white, acrid.



LACTARIUS (PIPERITES) FLEXUOSUS. *Fries.*
in pine woods, &c. Hayward Forest, 1881.

1700

2021

No 937

PL. 993.




LACTARIUS (PIPERITES) PYROGALUS. Bull.
in woods, fields, &c. Dinedor, 1881.

Mr. U

1000

100



LACTARIUS (PIPERITES) SCOTICUS. Berk. & Br.
amongst moss. Aboyne.

W 70 U

UoM

no. 434

PL. 977.

milk white, peppery.



LACTARIUS (PIPERITES) CAPSICUM. *Schulz.*
in grassy places. Dumfries, Sept. 1888.

1950

UofM

May 11



LACTARIUS (PIPERITES) CHRYSORHEUS. *Fries.*
in beech woods. Black Park, Langley, 1881.

11700

UofM .

W 70 U



LACTARIUS (PIPERITES) ACRIS. Bolton.
in woods. Reigate Hill.

W 70 U

Work

1871

No 9-10

PL 1006.

milk white, acrid.



LACTARIUS (PIPERITES) UMBRINUS. Pers.
in damp places. Epping Forest.

THE ...

THE ...

U of M

MrOU

PL 978.



LACTARIUS (PIPERITES) PARGAMENUS. *Swartz.*
on the ground. Hayward Forest, Sept. 1882.

Mr. U.S.

1074


milk copious, white, acrid.



LACTARIUS (PIPERITES) PIPERATUS. *Scop.*
on the ground in woods. Common.

Mr. U

vor!



No 945 |

PL. 980.



LACTARIUS (PIPERITES) VELLEREUS. *Fries.*
on ground in woods. Hayward Forest, 1883.

1800



1800

UoP 34


Mr. U



LACTARIUS (PIPERITES) EXSUCCUS. *Otto.*
on the ground. Epping Forest.



UoP M



no 447

PL. 982.



LACTARIUS (DAPETES) DELICIOSUS. Linn.
under firs. Lyndhurst, New Forest, 1882.

1000

44011

no. 975

PL. 1007.

milk mild, white.



LACTARIUS (RUSSULARIA) PALLIDUS. Pers.
in beech woods. Dinedor, 1881.

Wrot

UopA

no 444

PL. 983.



LACTARIUS (RUSSULARIA) QUIETUS. *Fries.*
in woods. Chingford, Essex. Nov. 1882.

1950



no. 950

PL. 1099.




LACTARIUS (*RUSSULARIA*) AURANTIACUS. *Fries.*
in woods. Woodman's Glade, Epping. Sept. 1888.

8700

UoM

no 721

PL. 1008.




LACTARIUS (RUSSULARIA) CREMOR. *Fries. var. PAUPER. Karst.*
in woods, &c. Carlisle, Sept. 1887.

And



UorN

Mr. U





#703



Uor M

2000





LACTARIUS (*RUSSULARIA*) CYATHULA. *Fries.*
under birches. *Minstead, New Forest.* *Aug. 1888.*

1000




Uor M

1901



LACTARIUS (*RUSSULARIA*) RUFUS. *Scop.*
in pine woods. near Abergelle.

1750



UoR

Mr. U




LACTARIUS (RUSSULARIA) HELVUS. *Fries.*
amongst moss in moist woods.

1701

Uer M

No 956

PL. 1010.



LACTARIUS (RUSSULARIA) TOMENTOSUS. Otto.
in damp places. Orton Moss, Carlisle, 1886.

W7011

U 74



LACTARIUS (RUSSULARIA) MAMMOSUS. var. MONSTROSUS. *Fries Icon.*
on the ground in woods. Scarborough, Oct. 1884.

1100

UorW



PL. 1011.

milk mild, white.



fragrant.

6-10 M

LACTARIUS (RUSSULARIA) GLYCIOUSMUS. *Fries.*
on the ground. Dinmore, Sept. 1888.


1701

U. 4

~~~~~


no 959

PL. 996.



LACTARIUS (RUSSULARIA) FULIGINOSUS. *Fries.*  
*in shady woods. Epping Forest, 1882.*

4700



1100

Mr. U

*milk white, acrid.*



**LACTARIUS (RUSSULARIA) PICINUS.** *Fries.*  
*in swampy ground. Foxley, 1879.*

MyoU

Uer 18

44 444





Roll

U of M

37 27

milk white, sweet.



LACTARIUS (RUSSULARIA) VOLEMUS. *Fries.*  
 in woods, &c. Breinton, Aug. 1873.

1700



UorM

1770

1770

1770

1770

1770



LACTARIUS (RUSSULARIA) ICHORATUS. Batsch.  
in woods, &c. near Hereford.

1901



UofM

M40U



*milk insipid, watery.*



**LACTARIUS (RUSSULARIA) SERIFLUUS.** *De Cand.*  
*in damp places. Haywood Forest, 1882.*

W701

Uor M

Mr. U





LACTARIUS (RUSSULARIA) MITISSIMUS. *Fries.*  
*in woods, &c. Dinedor Common, Oct. 1875.*

Mr. U



UofM

Mr. U



no 966



PL. 1002.

*milk white, mild.*



LACTARIUS (*RUSSULARIA*) SUBDULCIS. Bull.  
in woods. Epping Forest.

MyoU



U. of M.





odour fragrant when dried.  
milk mild, white.

A

DM

W.P.

LACTARIUS (RUSSULARIA) CAMPHORATUS. Bull.  
in woods. The Wrekin. Sept. 1875.

B

milk rather acrid, white.

odour of iugs.

LACTARIUS (RUSSULARIA) CENICARIUS. Batsch.  
amongst grass. Kew Gardens, 1882.

My 20

UofB



*milk watery, whitish, mild.*



M.

LACTARIUS (RUSSULARIA) SUBUMBONATUS. *Lind.*  
*in woods. Chingford, Epping Forest, 1886.*



LACTARIUS (RUSSULARIA) MINIMUS. *Smith.*  
*in fields, &c. Abergavenny.*

1700

UofM

M 70 U






LACTARIUS (RUSSULARIA) CENICARIUS. Batsch.  
amongst grass. Kew Gardens, 1882.

1701

UorA




*milk watery, whitish, mild.*



G.M.

**LACTARIUS (RUSSULARIA) SUBUMBONATUS.** *Lind.*  
*in woods. Chingford, Epping Forest, 1888.*



**LACTARIUS (RUSSULARIA) MINIMUS.** *Smith.*  
*in fields, &c. Abergavenny.*

UOF

May


1075

and



207

PL. 1014.



LACTARIUS (RUSSULARIA) OBNUBILIS. Lasch.  
in woods.

LACTARIUS (PLEUROPUS) OBLIQUUS. Fries.  
on banks. Dinmore.

1700

1700

1700

Uer 24




000000




RUSSULA (COMPACTÆ) NIGRICANS. *Bull.*  
*on the ground. Arboretum, Kew. July, 1888.*

May 11



UorM


Mr. O






*RUSSULA (COMPACTÆ) ALBO-NIGRA.* Kromb.  
amongst grass. Arboretum, Kew. Aug 1886.

WFOU



Uor M

M 30 U



turning black.

gills thin, crowded.

8 μ


RUSSULA (COMPACTÆ) ADUSTA. *Fries.*  
under trees. Minstead, New Forest. Aug. 1888.

8701



1074

1000



no 973

PL. 1017.



RUSSULA (COMPACTÆ) DENSIFOLIA. Gillet.  
under trees, amongst grass. Kew, Aug. 1886.

1702


UoF M





RUSSULA (COMPACTÆ) SEMICREMA. *Fries.*  
on the ground. *Neasden.* Sept. 1881.

M70U



no 712

PL. 1068.



RUSSULA (COMPACTÆ) DELICA. *Fries.*  
on the ground. *Kenilworth, and Neasden.* Oct. 1882.

Mr. U

UorM


1950

1950



**RUSSULA (COMPACTÆ) MUSTELINA.** *Frise.*  
*in woods. Slough, and Kew Gardens.*

1940




JOHN

2014




**RUSSULA (FURCATÆ) OLIVASCENS.** *Fries.*  
*on the ground.    Pleasure Grounds, Kew.    Aug. 1886.*

W 700



U.S. M.

2400





**RUSSULA (FURCATÆ) FURCATA. Fries.**  
*under trees. Fairmead, Epping, Aug. 1888.*

1700

UofM

Roll

\_\_\_\_\_




**RUSSULA (FURCATÆ) FURCATA. Fr. var. PICTIPES.**  
*under trees. Cuckoo Pits, Epping. Aug. 1888.*

1701



Uor M

Wrou



RUSSULA (*FURCATÆ*) FURCATA. *Fries, var. OCHROVIRIDIS.*  
*on the ground. Dinedor. Aug 1882.*

1901

UcrM

W750

No 931

PL. 1019.



*RUSSULA (FURCATÆ) SANGUINEA. Bull.*  
*on the ground. Haywood Common.*

Mr. U

UoP N

UoP N

Mr. U

No 982

PL. 1020.



RUSSULA (FURCATÆ) ROSACEA. *Fries.*  
amongst grass. *Credinhill Camp.* *Sept. 1882.*

1701

1702

Uor M

1400

acid.



RUSSULA (FURCATÆ) MACULATA. Quelet.  
under trees. Woodman's Glade, Epping. Sept. 1888.

Mr. U

Ver M

1970

No 984

PL. 1037.



*RUSSULA (FURCATÆ) SARDONIA. Fries. (variety).*  
*about larch. Black Park, Langley.*

4400

UoF M

87011



RUSSULA (*FURCATÆ*) DEPALLENS. *Pers.*  
*in woods. Dinedor, Sept. 1881.*

Mr. U



11



RUSSULA (FURCATÆ) PURPUREA. *Gillet.*  
amongst grass, &c. *Arboretum, Kew.*

1000

Uor M

W70U

PL. 1052.

*mild.*



RUSSULA (*FURCATÆ*) CÆRULEA. *Pers.*  
*amongst grass. Minstead, New Forest. Aug. 1888.*

WOL

UoP

1000



RUSSULA (*FURCATÆ*) DRIMEIA. Cooke.  
under trees. Black Park, Langley, 1881.

Wrol

1109

49011

PL. 1070.

*mild.*



*RUSSULA (RIGIDÆ) LACTEA. Pers.*  
*on the ground. Credinhill Camp. Sept. 1882.*

1901

UoP M

Mr. U

*mealy, mild.*



**RUSSULA (RIGIDÆ) LACTEA.** *Pers. var. INCARNATA. Quel.*  
*under pines, &c. Epping Forest.*


MrOU



200

26 991

PL. 1039.



**RUSSULA (RIGIDÆ) VIRESCENS. Schæff.**  
**under trees, Chingford, Epping Forest, Aug. 1884.**

Woll

U of W





RUSSULA (RIGIDÆ) CUTEFRACTA. Cooke.  
on the ground. Arboretum, Kew, 1881.

WroU

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

107 11

1000

No 993

PL. 1040.



RUSSULA (RIGIDÆ) CUTEFRACTA. Cooke.  
on the ground. Monk's Wood Epping 1881

Mr. U

Uor 34

Mrou

20994

PL. 1072.



RUSSULA (*RIGIDÆ*) LEPIDA. *Fries.*  
under trees. near Morpeth. Aug. 1888.

Mr. U

Uor M





RUSSULA (*RIGIDÆ*) LEPIDA. *Fries.* VARIETY.  
amongst grass. *Fairmead, Epping.* Aug. 1888.

Mr. U

11



*taste mild, sweet.*



**RUSSULA (RIGIDÆ) LINNÆI. Fries.**  
*on the ground. Crackley Wood, Kenilworth, 1885.*

11701

Uor M

Woll

Uor M

85

PL. 1073.

*mild.*



RUSSULA (*RIGIDÆ*) LEPIDA. *Fries.* VARIETY.  
*amongst grass. Fairmead, Epping. Aug. 1888.*

Mr. U

10





*RUSSULA (RIGIDÆ) LINNÆI. Fries.*  
*on the ground. Crackley Wood, Kenilworth, 1885.*

1961

UoP M

117



**RUSSULA (RIGIDÆ) XERAMPELINA.** *Schaff.*  
amongst grass. *Arboretum, Kew.*

8701



Woll

1701



*RUSSULA (RIGIDÆ) RUBRA. Fries.*  
*under trees. Monkswood, Epping.*



RUSSULA (RIGIDÆ) RUBRA. Fries. var. SAPIDA.  
under trees. Arboretum, Kew. Sept. 1888.

870U

Uor 34

W40U

PL. 1074.



RUSSULA (RIGIDA) XERAMPELINA. Schaaff,  
under trees. Arboretum, Kew. Sept. 1888.

Mr. U

UoM

1000

No. 1041

PL. 1041.



RUSSULA (RIGIDÆ) OLIVACEA. Schæff.  
amongst grass. Pleasure Grounds, Kew. Aug. 1888.

1901

Ugru

MrOU



RUSSULA (HETEROPHYLLÆ) VESCA. *Fries.*  
*amongst grass. Arboretum, Kew. 1886.*

Mr. U

U.S. 94





RUSSULA (RIGIDÆ) SEROTINA. *Quelet.*  
under beech trees. Arboretum, Kew. Sept. 1886.

Mr. U

Uor 14



PL 1054.

mild.



RUSSULA (HETEROPHYLLÆ) LILACEA. *Quelet.*  
*under trees. Loughton, Essex. Aug. 1881.*

Mr. U

11071

9789



RUSSULA (HETEROPHYLLÆ) AZUREA. *Bres.*  
*amongst grass. Arboretum, Kew, 1888.*


३१०॥

UrbM

1701

201006

PL. 1043.



**RUSSULA (HETEROPHYLLÆ) CYANOXANTHA.** *Schæff. (variety).*  
*amongst grass. Arboretum, Kew. Aug. 1888.*

0701

1074

WroU



RUSSULA (HETEROPHYLLÆ) CYANOXANTHA. *Schaff.*  
under trees. Arboretum, Kew. Aug. 1888.

Mr. U

UoR M

WFOU



RUSSULA (HETEROPHYLLÆ) CYANOXANTHA. Schaff. VARIETY.  
under trees. Arboretum, Kew. Sept. 1888.

Mr. U

UofM

100



**RUSSULA (HETEROPHYLLÆ) HETEROPHYLLA.** *Fries.* = *R. heterophylla*  
amongst grass. *Arboretum, Kew.* Aug. 1888.

450

UofM

---

1701



**RUSSULA (HETEROPHYLLÆ) HETEROPHYLLA.** *Fries.*  
*under beech. Arboretum, Kew. Aug. 1886.*

8700

Uay 𐄂𐄃𐄄

---

100

100



RUSSULA (HETEROPHYLLÆ) GALOCHROA. Bull.  
under trees. Arboretum, Kew. Aug. 1888.

M 70 U

U of M



Mr. U





*RUSSULA (HETEROPHYLLÆ) CONSOBRINA. Fries*  
*under beech. Kew Gardens. Aug. 1886.*

Mr. U



UoF M

1870

*very acrid.*

**RUSSULA (HETEROPHYLLÆ) CONSOBRINA. Fr. var. INTERMEDIA**  
*under trees. Arboretum, Kew. July, 1888.*



U of M

M7011

acid.



*RUSSULA (HETEROPHYLLÆ) CONSOBRINA. Fries. var. SORORIA.*  
amongst grass. Kew Gardens. July, 1884.

184011

Uop M

1901





**RUSSULA (HETEROPHYLLÆ) FOETENS.** *Pers.*  
*under trees. Black Park, Langley. Sept. 1881.*

M70U

UofM



Mr. C.



*somewhat acrid,  
odour unpleasant.*



10-8 M

*whole plant tough and elastic.*

**RUSSULA (HETEROPHYLLÆ) SUBCÆTENS.** *Smith.*  
*amongst grass. Pleasure Grounds, Kew. Aug. 1886.*



Mr O



form





RUSSULA (HETEROPHYLLÆ) FELLEA. *Fries.*  
*under trees. Arboretum, Kew. Sept. 1881.*



M76U

UoFM





**RUSSULA (HETEROPHYLLÆ) ELEGANS.** *Bresadola.*  
*amongst grass. Arboretum, Kew. Sept. 1886.*

1870

U of M

1900



*RUSSULA (HETEROPHYLLÆ) QUELETII. Fries.*  
*under firs. Black Park, Langley.*

M76U

UorM





*RUSSULA (HETEROPHYLLÆ) EXPALLENS. Gillet.*  
*under trees. Arboretum, Kew, Oct. 1886.*

M70U



UofM

THE UNIVERSITY OF MICHIGAN LIBRARY  
ANN ARBOR, MICHIGAN 48106-1069

W 7011



RUSSULA (*FRAGILES*) EMETICA. *Fries.*  
under trees in woods. *Neasden, Sept. 1881.*

M70U

1278

W. 1. 11



**RUSSULA (FRAGILES) CLUSII.** *Fries.*  
*under trees. (A) Epping Forest. (B) Black Park, 1881.*



Uor M



Mr. U



No 1023

PL 1059.



RUSSULA (FRAGILES) FALLAX Schaff.  
on the ground. Neasden. Sept. 1881.

Mr. U

UorM






*slightly acid.*




**RUSSULA (FRAGILES) PECTINATA. Bull.**  
*on the ground. Breinton, Hereford, 1881.*



U or M







**RUSSULA (FRAGILES) OCHROLEUCA.** *Pers.*  
*on the ground. Arboretum, Kew, 1881.*

Who U



UoRM





*RUSSULA (FURCATÆ) GRANULOSA.* Cooke.  
under trees. Arboretum, Kew. Aug. 1888.



UorM

1870



mild.



RUSSULA (FRAGILES) AERUGINEA, Fries.  
on the ground. King's Langley.



M70U




*Boletus fragilis* (Fr.)  
on the ground. Drying Press.

Uor M



acrid.




RUSSULA (FRAGILES) FRAGILIS. Fries.  
on the ground. Epping Forest.

Mr. U

UorM







W 4011



**RUSSULA (FRAGILES) FINGIBILIS.** *Britz.*  
*under trees. Arboretum, Kew. July, 1882.*




M70U

UorM



1170



mild.



RUSSULA (FRAGILES) FINGIBILIS. Britz.  
under trees. Arboretum, Kew. July, 1882.

Mr. U

Uor M





*mild, then slightly acrid.*



RUSSULA (FRAGILES) CITRINA. Gillet.  
on ground in woods. Black Park. Sept. 1881.

84 no 11






*RUSSULA (FRAGILES) PUNCTATA. Gillet, var. LEUCOPUS.*  
*amongst grass. Arboretum, Kew. July, 1888.*

Uro



Uor M







RUSSULA (*FRAGILES*) VETERIOSA. *Fries.*  
amongst grass under trees. *Pleasure Grounds, Kew, 1886.*

1790



Uor M







RUSSULA (*FRAGILES*) VETERIOSA. *Fries.*  
*under trees. Arboretum, Kew. Sept. 1888.*

Mr. U



Uor M



Whole



RUSSULA (*FRAGILES*) ROSEIPES. *Secr. VARIETY.*  
amongst grass. *Morpeth. Sept. 1888.*

1040


Uor M



18-11-11




*viscid, mild.*



*RUSSULA (FRAGILES) INTEGRA. Linn.*  
*under trees. Arboretum, Kew.*

M70U

UoFM





RUSSULA (FRAGILES) INTEGRA. Linn.  
on the ground. Fairmead, Epping, 1882.

1901

UoM





RUSSULA (FRAGILES) INTEGRA. Linn. var. ALBA.  
under trees. Sutton Park, Warwickshire.



U of M





*mild.*




**RUSSULA (FRAGILES) DECOLORANS.** *Fries.*  
*in pine woods. Aboyme, 1872.*



4400

UorM





no 1040

PL. 1061.




RUSSULA (FRAGILES) BARLÆ. Quelet.  
amongst grass. Arboretum, Kew. July, 1888.

100-1000

U of M







RUSSULA (*FRAGILES*) AURATA. *With.*  
*in woods. Horsham.*

RhoU

U of M



.....





*RUSSULA (FRAGILES) NITIDA. Fries. var. B Secr. 496.*  
*in woods. Carlisle. Aug. 1888.*


1870



1000

Mholi





*RUSSULA (FRAGILES) NITIDA. Pers.*  
*in woods. (A) Laxton, Norths. (B) Arboreium, Kew.*


Myou

U or M



\_\_\_\_\_

22





RUSSULA (*FRAGILES*) NITIDA. *Fr. var. CUPRÆA. Krom.*  
under firs. Bramham Park, Leeds, 1888.

WFOU

UorM



1111

*very fragile.*

**RUSSULA (FRAGILES) ARMENIACA.** *Cooke.*  
*amongst grass, under trees. Fairmead, Epping. Aug. 1888.*




M70U



U of M





20176

PL. 1065.

*mild.*



RUSSULA (*FRAGILES*) PUELLARIS. *Fries.*  
*under birch. Morpetk, Sept. 1888.*

8700

U of M



Mr. U. S. S.



no 1047

PL. 1068.



turning fovey where touched.

**RUSSULA (FRAGILES) PUELLARIS. Fries. VARIETY.**  
*under trees in woods. Morpeth, Sept. 1888.*

1901



UofM



Mr. U



RUSSULA (*FRAGILES*) ALUTACEA. *Fries.*  
amongst trees. *Minstead, New Forest. Aug. 1888.*

WFOU



UoF M





No 1049

PL. 1097.

mild



RUSSULA (*FRAGILES*) ALUTACEA. *Fries.*  
*on the ground. Epping Forest, 1881.*

Mr. U.




U or M



1810






*RUSSULA (FRAGILES) OCHRACEA. Alb. & Schw.*  
*amongst grass. Pleasure Grounds, Kew. Aug. 1886.*



117011

Uor M








RUSSULA (*FRAGILES*) LUTEA. Hudson.  
in woody places. Hayward Forest, 1881.

Mr. U



UoP





401052

PL. 1147.




**RUSSULA (FRAGILES) NAUSEOSA.** *Fries.*  
*on the ground. Morpeth, 1889.*

0400

UoF M







RUSSULA (FRAGILES) VITELLINA. Fries.  
on the ground. Chingford, Essex. Aug. 1886.



Mr. H. U.

UoP M



1963






**RUSSULA (FRAGILES) CHAMELEONTINA.** *Fries.*  
*under beeches. Arboretum, Kew, 1886.*

1901

Uor M





**CANTHARELLUS CIBARIUS.** *Fries.*  
*on the ground. Epping Forest, 1886.*

WFOU

Uor M







A

CANTHARELLUS CIBARIUS. Fr. var. RUFIPES. Gill.  
on the ground. Reading, Oct. 1872.



B

tomentose.

CANTHARELLUS FRIESII. Quelst.  
in woods. Sydenham.




no 1007

PL. 1104.



CANTHARELLUS AURANTIACUS. Fries.  
amongst grass. Forest of Dean, 1887.



M70U



UoP M





CANTHARELLUS BROWNII. B. & Br.  
*in grassy places. Hitchin.*

CANTHARELLUS UMBONATUS. Fries.  
*amongst moss. Mossburnford.*

1870



U of M



1890




CANTHARELLUS CARBONARIUS. *Fries.*  
 on charred ground. *Arboretum, Kew.* Nov. 1880.




1870

Uor M








**CANTHARELLUS HOUGHTONI.** *Phillips.*  
on the ground, near Hereford.


Mr. U




U of M







CANTHARELLUS TUBÆFORMIS. *Fries.*  
on the ground. *King's Cliffe.*



1904

U of M







CANTHARELLUS INFUNDIBULIFORMIS. *Fries.*  
*on the ground. Carlisle, Oct. 1887.*

Mr. U

U of M





116  
(a)



CANTHARELLUS CINEREUS. Fries.  
on the ground. Cord Cuck.



CANTHARELLUS CUPULATUS. Fries.  
on the ground. near Bristol. Nov. 1882.


C.B.

7 x 5 μ

116

U90M

Uor M





A



3 X 5 M

CANTHARELLUS LEUCOPHAeus. *Novel.*  
*on the ground. King's Cliffs.*



B

CANTHARELLUS STEVENSONI. *B. & Br.*  
*on stumps. Glamis, N.B.*

1701

UorM





A



CANTHARELLUS MUSCIGENUS. *Bull.*  
on mosses. *Aboyne.*

B



W.P.

UorM

CANTHARELLUS GLAUCUS. *Batsch.*  
on sandy slopes. *near Shrewsbury.*

M70U

UoFM





CANTHARELLUS RETIRUGUS. *Fries.*  
on mosses. *King's Cliffe.*

CANTHARELLUS LOBATUS. *Fries.*  
in swamps. *Apetkorp.*

Mr. U




Uor B



1700





NYCTALIS CALIGINOSA. Smith.  
amongst leaves. Bishop's Wood, Highgate.

NYCTALIS ASTEROPHORA. Fries.  
on dead *Russula*. Epping Forest.

M70U

Ua



No 1068

PL. 1113.




NYCTALIS PARASITICA. Fries.  
on decaying fungi. Hainault Forest.

Mr. U

UofM





No 1869

PL. 1118.




MARASMIUS (SCORTEI) URENS. Bull.  
amongst grass. Pleasure Grounds, Kew. Sept. 1884.

M70U

Uor M








MARASMIUS (SCORTEI) PERONATUS. Fries.  
*on the ground. Hainault Forest.*






UFOU


Uor M.



odour of garlic.



MARASMIUS (SCORTEI) PORREUS. Pers.  
amongst dead leaves. Collyweston.



1704

Uor 3






No 1072

PL. 1118.



MARASMIUS (COLLYBIA) OREADES. *Fries.*  
on lawns, &c. *Kew Gardens.*



M76U


12

12

Uor M



1871





**MARASMIUS (SCORTEI) SCORTEUS.** *Fries.*  
*in moist woods. Laxton.*

M 70 U

U of M





no 1074

PL. 1120.

*odour of garlic.*



MARASMIUS (TERBINI) PRASIOSMUS. Fries.  
amongst leaves. Scarborough.

M70U

UoM



W. H. O. U.





MARASMIUS (TERGINI) VARICOSUS. Fries.  
in moist places, amongst moss.



MARASMIUS (TERGINI) FUSCO-PURPUREUS. Pers.  
amongst twigs and leaves. Apethorpe.

M70U

UorM







MARASMIUS (TERBINI) TERGINUS. Fries.  
amongst leaves. near Bristol.




MARASMIUS (TERBINI) ARCHYROPUS. Fries.  
amongst leaves. near Bristol.

M70U



Uor M








MARASMIUS (TERBINI) ERYTHROPUS. Pers.  
amongst leaves. Fineshade, Notts.

M70U

Uor M








MARASMIUS (TERGINI) IMPUDICUS. Fries.  
about fir trees. Hanham, Wilts.

ou

Uor M





*strong scented.*

A



*stem smooth.*

MARASMIUS (CALOPODES) SCORODONIUS. *Pries.*  
*on twigs, &c. near Hereford.*

B



*7 x 4 M*

*stem smooth.*


MARASMIUS (CALOPODES) CALOPUS. *Pers.*  
*on twigs, &c. near Bristol.*

1096


Work








MARASMIUS (CALOPODES) VAILLANTII. *Fries.*  
on twigs and leaves.



MARASMIUS (CALOPODES) ANGULATUS. *Pers.*  
on submerged rushes. *Exeter, Aug 1884.*



MARASMIUS (CALOPODES) LANGUIDUS. *Lasch.*  
on twigs, &c. *Frome.*



1070 M

UOPM



M 40 U





MARASMIUS (CALOPODES) FOETIDUS. Sow  
on branches. Pontrilas, 1885.

MARASMIUS (CHORDALES) CAUTICINALIS. With.  
amongst pine leaves, &c. near Hereford.



1700







MARASMIUS AMADELPHUS. Bull.  
on twigs. Bath.

MARASMIUS CANDIDUS. Fries.  
after Bolton.

MARASMIUS (CALOPODES) RAMEALIS. Bull.  
on bramble, &c. Chingford, Essex. Oct. 1882.

M70U

UofM





MARASMIUS (MYCENA) COHÆRENS. Fries.  
on the ground. Downton.

Mr. U

Uorh






N.J.B.

stem smooth.


6 X 3-4 M

MARASMIUS (ROTULÆ) ROTULA. Scop.  
on dead twigs. King's Cliffs.



stem smooth.


MARASMIUS (ROTULÆ) GRAMINUM. Lab.  
on grass, &c. Fineshade, Norfolk.




7 X 3-4 M

stem smooth.

MARASMIUS (ROTULÆ) ANDROSACEUS. Bull.  
amongst leaves. Epping Forest.



UorM





MARASMIUS (ROTULÆ) SPLACHNOIDES. Fries.  
among pine leaves. Calatya.



MARASMIUS (ROTULÆ) CURREYI. B. & Br.  
on grass. Finckhade, Notts.



MARASMIUS (ROTULÆ) PERFORANS. Fries.  
on larch leaves. Coed Coch.

1 4 7 7


1 4 7 7


1 1 4 7 7

1700

Uor M








MARASMIUS (*ROTULÆ*) HUDSONI. Pers.  
on dead holly leaves. Highgate.

FORM

Uor 34







Mr. U


No 1088







MARASMIUS (ROTULÆ) EPIPHYLLUS. Fries.  
on dead leaves. Epping.



MARASMIUS (ROTULÆ) POLYADELPHUS. Lenz.  
on dead leaves, &c. Dinmore



= *M. Broomei*. B.

MARASMIUS (APUS) SPODOLEUCUS. B. & Br.  
on elm twigs. Batheaston.


U90M

Uer 81





LENTINUS TIGRINUS. *Fries.*  
*on fir in greenhouse. King's Lynn. Aug. 1884.*



Uor 14








H.G.B.

LENTINUS TIGRINUS. *Fries.*  
on wood. *Reigate, July, 1875.*




M.J.B.

LENTINUS DUNALII. *Fries.*  
on ash. *Dorsetshire.*



Uop 17





251091


PL. 1140.



LENTINUS LEPEDEUS. *Fries.*  
on dead wood. *Mattishall, Norfolk. May, 1881.*


U90M

Upp 88





*LENTINUS LEPIDEUS. Fries. (monstrous form.)*  
*on wood. Brewery. Burton. Mar. 1883.*






M70 U




184




1040










WFOU







PANUS STYPTICUS. *Fries.*  
on stumps. *Downton, 1889.*




PANUS FARINACEUS. *Fries.*  
on branches. *Epping Forest.*



PANUS PATELLARIS. *Fries.*  
on cherry. *Forres, N.B.*

1903

100177






XEROTUS DEGENER.  
on the naked ground. Fries.

1701




U or M





TROGIA CRISPA. Fries.  
on branches. Morpeth. Dec. 1883.





SCHIZOPHYLLUM COMMUNE. Fries.  
on logs. near Ely.

1904



UorN





LENZITES BETULINA. *Fries.*  
on stumps. *Epping Forest.*

LENZITES FLACCIDA. *Bull.*  
on beech. *Hainault Forest.*

WOL

Mr. U



Uor M



LENZITES SEPIARIA Fries.  
on fir trunks. Northamptonshire.


B

7-8 x 4M


LENZITES ABIETINA Fries.  
on fir trunks and branches. Glasgow.

W40U









1871







BOUND

MAY 9 1968

UNIV. OF MICH.  
LIBRARY

1.

